

## 1. Dressing Care:

Change your dressing two days after surgery. Some light bleeding is normal. Showering is permitted 2 days after surgery & the incision may be washed with soap & water, but no submerging or bathing is allowed until 2 weeks. Keep a clean dressing on the incision until your follow-up appointment.

## 2. Medication:

You will be given a painkiller \_\_\_\_\_ to be taken 1-2 pills every 4-6 hours as needed. Do not drink or drive while on pain medication. Do not take additional Tylenol based pain medications while on this medication.

Take calcium with vitamin D supplements 500mg 3x day with meals.

Do not take any anti-inflammatory medication as this may impede bone healing (ibuprofen, Motrin, Naprosyn, Celebrex, etc.).

## 3. Activity:

You will be placed in a sling after surgery but should remove the arm from the sling to work on gentle passive pendulum range of motion exercises daily as instructed. No other motion or activity is allowed at this time as this could result in a failure of the repair with displacement of the fracture.

Motion may be gradually increased as pain allows at 2 weeks. Therapy will be started at 8 weeks to work on restoring full motion and rotator cuff strengthening, progressing to full activity as tolerated. Expected time to complete recovery is 6 months.

## 4. General Instructions:

Do not smoke! Nicotine impedes bone healing and can prevent or delay fracture union.

If you notice any signs of infection such as increased pain, redness, wound drainage or fever  $>100.5^{\circ}$  F, call the office immediately.

## 5. Follow-up Appointment:

Call the office at (616) 949-8945 to schedule an appointment for 10-14 days after surgery.

## 6. Other Instructions:

---

*If you have any problems or concerns following surgery,  
please call our office at (616) 949-8945.*