

1. Splint Care:

Keep your splint clean & dry. Do not remove. Swelling is normal and may persist for several weeks. If the splint feels too tight, ice & elevate the leg to reduce swelling. If you have progressive pain or numbness in the leg, call the office.

2. Medication:

You will be given a painkiller _____ to be taken 1-2 pills every 4-6 hours as needed. Do not drink or drive while on pain medication. Do not take additional Tylenol based pain medications while on this medication.

Take an aspirin 325mg once a day to help prevent blood clots unless you are already on a blood thinning medication.

3. Activity:

No weight bearing is allowed on the affected leg for 8-12 weeks. Any walking or weight bearing even in the splint or cast could result in a failure of the repair with displacement of the fracture. You may begin walking at 8-12 weeks once X-Rays have shown the fracture to be healed.

Expected time to complete recovery is 4-6 months.

4. General Instructions:

Apply ice to the splint over the ankle (20min on /off) and elevate to reduce pain & swelling.

If you notice any signs of infection such as increased pain, redness, wound drainage or fever $>100.5^{\circ}$ F, call the office immediately.

If you develop any calf or thigh pain, increased swelling, chest pain or shortness of breath, call the office immediately or go to the ER as these are symptoms of a deep venous thrombosis (blood clot) in the leg.

5. Follow-up Appointment:

Call the office at (616) 949-8945 to schedule an appointment for 10-14 days after surgery.

6. Other Instructions:

*If you have any problems or concerns following surgery, please call our office at
(616) 949-8945.*